

Pediatric Associates of Lagrange
205 Calumet Center Road
Lagrange, GA 30241
(706) 885-1961

Lisa P. Allardice

Congratulations on your new bundle of joy. Oftentimes along with joy come feelings of sadness or being overwhelmed. Our office cares about your family and wants you to know we are here for you. It is recommended by the American Academy of Pediatrics to screen parents for postpartum depression. Our office wants to be supportive of efforts to address postpartum issues. We will ask you to complete this screening tool at newborn through your baby's first year visits. Depending on your score, we have resources that we can offer you if you have any signs of postpartum depression or just need to talk to someone about what you're feeling.

Postpartum depression is not the same as baby blues, which you may be familiar with. Postpartum depression lasts longer and is more serious than baby blues. Baby blues are feelings of sadness you may have in the first few days after having a baby. Baby blues can happen 2 to 3 days after you have your baby and can last up to 2 weeks. You may have trouble sleeping, be moody or cranky, and cry a lot. Postpartum depression (also called PPD) is a kind of depression that some women get after having a baby. It is strong feelings of sadness, anxiety (worry) and tiredness that last for a long time after giving birth. These feelings can make it hard for you to take care of yourself and your baby. PPD can happen any time after childbirth, but it often starts within 1 to 3 weeks of having a baby. Please remember PPD is not your fault and you didn't do anything to cause PPD. It doesn't make you a bad person or a bad mother. You are not alone. Please feel free to reach out to us since up to 1 in 7 women has PPD after giving birth, and it's the most common complication for women who have just had a baby. It's a medical condition that needs treatment to get better. If you have feelings that last longer than 2 weeks or if something doesn't seem right, call your healthcare provider.

With your permission we will share your scores with your OB/GYN. If needed, we will also give you information regarding resources and support networks.

- Permission to share with my OB/GYN or Primary Care Provider:

Name: _____ Phone: _____

- My contact information may be shared with referral sources:

Referral Source Name: _____ Phone: _____

Signature: _____ Date: _____

Print your name: _____

Edinburgh Postnatal Depression Scale¹ (EPDS)

Name: _____

Address: _____

Your Date of Birth: _____

Baby's Date of Birth: _____

Phone: _____

As you are pregnant or have recently had a baby, we would like to know how you are feeling. Please check the answer that comes closest to how you have felt **IN THE PAST 7 DAYS**, not just how you feel today.

Here is an example, already completed.

I have felt happy:

- Yes, all the time
- Yes, most of the time This would mean: "I have felt happy most of the time" during the past week.
- No, not very often Please complete the other questions in the same way.
- No, not at all

In the past 7 days:

1. I have been able to laugh and see the funny side of things
 - As much as I always could
 - Not quite so much now
 - Definitely not so much now
 - Not at all
2. I have looked forward with enjoyment to things
 - As much as I ever did
 - Rather less than I used to
 - Definitely less than I used to
 - Hardly at all
3. I have blamed myself unnecessarily when things went wrong
 - Yes, most of the time
 - Yes, some of the time
 - Not very often
 - No, never
4. I have been anxious or worried for no good reason
 - No, not at all
 - Hardly ever
 - Yes, sometimes
 - Yes, very often
5. I have felt scared or panicky for no very good reason
 - Yes, quite a lot
 - Yes, sometimes
 - No, not much
 - No, not at all
6. Things have been getting on top of me
 - Yes, most of the time I haven't been able to cope at all
 - Yes, sometimes I haven't been coping as well as usual
 - No, most of the time I have coped quite well
 - No, I have been coping as well as ever
7. I have been so unhappy that I have had difficulty sleeping
 - Yes, most of the time
 - Yes, sometimes
 - Not very often
 - No, not at all
8. I have felt sad or miserable
 - Yes, most of the time
 - Yes, quite often
 - Not very often
 - No, not at all
9. I have been so unhappy that I have been crying
 - Yes, most of the time
 - Yes, quite often
 - Only occasionally
 - No, never
10. The thought of harming myself has occurred to me
 - Yes, quite often
 - Sometimes
 - Hardly ever
 - Never

Administered/Reviewed by _____ Date _____

¹Source: Cox, J.L., Holden, J.M., and Sagovsky, R. 1987. Detection of postnatal depression: Development of the 10-item Edinburgh Postnatal Depression Scale. *British Journal of Psychiatry* 150:782-786 .

²Source: K. L. Wisner, B. L. Parry, C. M. Plontek, Postpartum Depression N Engl J Med vol. 347, No 3, July 18, 2002, 194-199

Users may reproduce the scale without further permission providing they respect copyright by quoting the names of the authors, the title and the source of the paper in all reproduced copies.

Patient Name: _____

Lead Risk Assessment Questionnaire

Circle Yes or No

*If Yes, please explain

- | | | | |
|------------------------------------------------------------------------------------------------------------------------|-----|----|-------|
| 1. Does your child live in or often visit a house that may have been built before 1978? | Yes | No | _____ |
| 2. Does your child live in or often visit a house that is being remodeled or is having paint removed? | Yes | No | _____ |
| 3. Does your child live with or often visit another child that has an elevated blood lead level? | Yes | No | _____ |
| 4. Does your child live with anyone that works at a job where lead may be found or has a hobby that uses it? | Yes | No | _____ |
| 5. Does your child chew on or eat any non-food items like paint chips or dirt? | Yes | No | _____ |
| 6. Does your child live near an active lead smelter, battery recycling plant or other industry likely to release lead? | Yes | No | _____ |
| 7. Does your child receive medicines such as <i>greta</i> , <i>azarcon</i> , <i>kohl</i> or <i>pay-loo-ah</i> ? | Yes | No | _____ |

Risk Factors for Hearing Loss

- | | | |
|-------------------------------------------------------------------------------------------------------|-----|----|
| 1. Parent or caregiver concern regarding hearing, speech, language, or developmental delay | Yes | No |
| 2. Family history of permanent childhood hearing loss | Yes | No |
| 3. NICU stay greater than five days, ECMO, ototoxic medications, loop diuretics, exchange transfusion | Yes | No |
| 4. In utero infections such as CMV, herpes, rubella, syphilis, Toxoplasmosis | Yes | No |
| 5. Craniofacial anomalies, especially involving the ear and temporal bone | Yes | No |
| 6. Stigmata of syndromes known to cause hearing loss, Eustachian tube dysfunction | Yes | No |
| 7. Syndromes associated with hearing loss or progressive or late-onset hearing loss | Yes | No |
| 8. Neurodegenerative disorders or sensory motor neuropathies | Yes | No |
| 9. Postnatal infections associated with sensorineural hearing loss including bacterial meningitis | Yes | No |
| 10. Head trauma | Yes | No |
| 11. Chemotherapy | Yes | No |
| 12. Recurrent or persistent ear infection for at least 3 months | Yes | No |

Parent Signature _____

Date _____

Please check the box and sign if there have been no changes to the above answers since the last checkup.

- | | | |
|-------------------------------------------------|---------------|------------|
| <input type="checkbox"/> Parent Signature _____ | Witness _____ | Date _____ |
| <input type="checkbox"/> Parent Signature _____ | Witness _____ | Date _____ |
| <input type="checkbox"/> Parent Signature _____ | Witness _____ | Date _____ |
| <input type="checkbox"/> Parent Signature _____ | Witness _____ | Date _____ |
| <input type="checkbox"/> Parent Signature _____ | Witness _____ | Date _____ |
| <input type="checkbox"/> Parent Signature _____ | Witness _____ | Date _____ |
| <input type="checkbox"/> Parent Signature _____ | Witness _____ | Date _____ |
| <input type="checkbox"/> Parent Signature _____ | Witness _____ | Date _____ |

Patient Name: _____

Hepatitis B Vaccine:

What You Need to Know

Many Vaccine Information Statements are available in Spanish and other languages. See www.immunize.org/vis

Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite www.immunize.org/vis

1 Why get vaccinated?

Hepatitis B vaccine can prevent **hepatitis B**. Hepatitis B is a liver disease that can cause mild illness lasting a few weeks, or it can lead to a serious, lifelong illness.

- **Acute hepatitis B infection** is a short-term illness that can lead to fever, fatigue, loss of appetite, nausea, vomiting, jaundice (yellow skin or eyes, dark urine, clay-colored bowel movements), and pain in the muscles, joints, and stomach.
- **Chronic hepatitis B infection** is a long-term illness that occurs when the hepatitis B virus remains in a person's body. Most people who go on to develop chronic hepatitis B do not have symptoms, but it is still very serious and can lead to liver damage (cirrhosis), liver cancer, and death. Chronically-infected people can spread hepatitis B virus to others, even if they do not feel or look sick themselves.

Hepatitis B is spread when blood, semen, or other body fluid infected with the hepatitis B virus enters the body of a person who is not infected. People can become infected through:

- Birth (if a mother has hepatitis B, her baby can become infected)
- Sharing items such as razors or toothbrushes with an infected person
- Contact with the blood or open sores of an infected person
- Sex with an infected partner
- Sharing needles, syringes, or other drug-injection equipment
- Exposure to blood from needlesticks or other sharp instruments

Most people who are vaccinated with hepatitis B vaccine are immune for life.

2 Hepatitis B vaccine

Hepatitis B vaccine is usually given as 2, 3, or 4 shots.

Infants should get their first dose of hepatitis B vaccine at birth and will usually complete the series at 6 months of age (sometimes it will take longer than 6 months to complete the series).

Children and adolescents younger than 19 years of age who have not yet gotten the vaccine should also be vaccinated.

Hepatitis B vaccine is also recommended for certain **unvaccinated adults**:

- People whose sex partners have hepatitis B
- Sexually active persons who are not in a long-term monogamous relationship
- Persons seeking evaluation or treatment for a sexually transmitted disease
- Men who have sexual contact with other men
- People who share needles, syringes, or other drug-injection equipment
- People who have household contact with someone infected with the hepatitis B virus
- Health care and public safety workers at risk for exposure to blood or body fluids
- Residents and staff of facilities for developmentally disabled persons
- Persons in correctional facilities
- Victims of sexual assault or abuse
- Travelers to regions with increased rates of hepatitis B
- People with chronic liver disease, kidney disease, HIV infection, infection with hepatitis C, or diabetes
- Anyone who wants to be protected from hepatitis B

Hepatitis B vaccine may be given at the same time as other vaccines.



3

Talk with your health care provider

Tell your vaccine provider if the person getting the vaccine:

- Has had an **allergic reaction after a previous dose of hepatitis B vaccine**, or has any **severe, life-threatening allergies**.

In some cases, your health care provider may decide to postpone hepatitis B vaccination to a future visit.

People with minor illnesses, such as a cold, may be vaccinated. People who are moderately or severely ill should usually wait until they recover before getting hepatitis B vaccine.

Your health care provider can give you more information.

4

Risks of a vaccine reaction

- Soreness where the shot is given or fever can happen after hepatitis B vaccine.

People sometimes faint after medical procedures, including vaccination. Tell your provider if you feel dizzy or have vision changes or ringing in the ears.

As with any medicine, there is a very remote chance of a vaccine causing a severe allergic reaction, other serious injury, or death.

5

What if there is a serious problem?

An allergic reaction could occur after the vaccinated person leaves the clinic. If you see signs of a severe allergic reaction (hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, or weakness), call **9-1-1** and get the person to the nearest hospital.

For other signs that concern you, call your health care provider.

Adverse reactions should be reported to the Vaccine Adverse Event Reporting System (VAERS). Your health care provider will usually file this report, or you can do it yourself. Visit the VAERS website at www.vaers.hhs.gov or call **1-800-822-7967**. *VAERS is only for reporting reactions, and VAERS staff do not give medical advice.*

6

The National Vaccine Injury Compensation Program

The National Vaccine Injury Compensation Program (VICP) is a federal program that was created to compensate people who may have been injured by certain vaccines. Visit the VICP website at www.hrsa.gov/vaccinecompensation or call **1-800-338-2382** to learn about the program and about filing a claim. There is a time limit to file a claim for compensation.

7

How can I learn more?

- Ask your healthcare provider.
- Call your local or state health department.
- Contact the Centers for Disease Control and Prevention (CDC):
 - Call **1-800-232-4636 (1-800-CDC-INFO)** or
 - Visit CDC's www.cdc.gov/vaccines

Vaccine Information Statement (Interim)
Hepatitis B Vaccine



Office use only

8/15/2019 | 42 U.S.C. § 300aa-26

Your Child's First Vaccines

What You Need to Know

Many Vaccine Information Statements are available in Spanish and other languages. See www.immunize.org/vis

Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite www.immunize.org/vis

The vaccines covered on this statement are those most likely to be given during the same visits during infancy and early childhood. Other vaccines (including measles, mumps, and rubella; varicella; rotavirus; influenza; and hepatitis A) are also routinely recommended during the first five years of life.

Your child will get these vaccines today:

DTaP Hib Hepatitis B Polio PCV13

(Provider: Check appropriate boxes.)

1 Why get vaccinated?

Vaccine-preventable diseases are much less common than they used to be, thanks to vaccination. But they have not gone away. Outbreaks of some of these diseases still occur across the United States. **When fewer babies get vaccinated, more babies get sick.**

7 childhood diseases that can be prevented by vaccines:

1. Diphtheria (the 'D' in DTaP vaccine)

- **Signs and symptoms** include a thick coating in the back of the throat that can make it hard to breathe.
- **Diphtheria can lead to** breathing problems, paralysis and heart failure.
 - About 15,000 people died each year in the U.S. from diphtheria before there was a vaccine.

2. Tetanus (the 'T' in DTaP vaccine; also known as Lockjaw)

- **Signs and symptoms** include painful tightening of the muscles, usually all over the body.
- **Tetanus can lead to** stiffness of the jaw that can make it difficult to open the mouth or swallow.
 - Tetanus kills about 1 person out of every 10 who get it.

3. Pertussis (the 'P' in DTaP vaccine, also known as Whooping Cough)

- **Signs and symptoms** include violent coughing spells that can make it hard for a baby to eat, drink, or breathe. These spells can last for several weeks.
- **Pertussis can lead to** pneumonia, seizures, brain damage, or death. Pertussis can be very dangerous in infants.
 - Most pertussis deaths are in babies younger than 3 months of age.

4. Hib (*Haemophilus influenzae* type b)

- **Signs and symptoms** can include fever, headache, stiff neck, cough, and shortness of breath. There might not be any signs or symptoms in mild cases.
- **Hib can lead to** meningitis (infection of the brain and spinal cord coverings); pneumonia; infections of the ears, sinuses, blood, joints, bones, and covering of the heart; brain damage; severe swelling of the throat, making it hard to breathe; and deafness.
 - Children younger than 5 years of age are at greatest risk for Hib disease.

5. Hepatitis B

- **Signs and symptoms** include tiredness, diarrhea and vomiting, jaundice (yellow skin or eyes), and pain in muscles, joints and stomach. But usually there are no signs or symptoms at all.
- **Hepatitis B can lead to** liver damage, and liver cancer. Some people develop chronic (long term) hepatitis B infection. These people might not look or feel sick, but they can infect others.
 - Hepatitis B can cause liver damage and cancer in 1 child out of 4 who are chronically infected.

6. Polio

- **Signs and symptoms** can include flu-like illness, or there may be no signs or symptoms at all.
- **Polio can lead to** permanent paralysis (can't move an arm or leg, or sometimes can't breathe) and death.
 - In the 1950s, polio paralyzed more than 15,000 people every year in the U.S.



7. Pneumococcal Disease

- **Signs and symptoms** include fever, chills, cough, and chest pain. In infants, symptoms can also include meningitis, seizures, and sometimes rash.
- **Pneumococcal disease can lead to** meningitis (infection of the brain and spinal cord coverings); infections of the ears, sinuses and blood; pneumonia; deafness; and brain damage.
 - About 1 out of 15 children who get pneumococcal meningitis will die from the infection.

Children usually catch these diseases from other children or adults, who might not even know they are infected. A mother infected with hepatitis B can infect her baby at birth. Tetanus enters the body through a cut or wound; it is not spread from person to person.

Vaccines that protect your baby from these seven diseases:

Vaccine	Number of doses	Recommended ages	Other information
DTaP (Diphtheria, Tetanus, Pertussis)	5	2 months, 4 months, 6 months, 15-18 months, 4-6 years	Some children get a vaccine called DT (Diphtheria & Tetanus) instead of DTaP.
Hepatitis B	3	Birth, 1-2 months, 6-18 months	
Polio	4	2 months, 4 months, 6-18 months, 4-6 years	An additional dose of polio vaccine may be recommended for travel to certain countries.
Hib (<i>Haemophilus influenzae</i> type b)	3 or 4	2 months, 4 months, (6 months), 12-15 months	There are several Hib vaccines. With one of them the 6-month dose is not needed.
Pneumococcal (PCV13)	4	2 months, 4 months, 6 months, 12-15 months	Older children with certain health conditions also need this vaccine.

Your healthcare provider might offer some of these vaccines as **combination vaccines**—several vaccines given in the same shot. Combination vaccines are as safe and effective as the individual vaccines, and can mean fewer shots for your baby.

2 Some children should not get certain vaccines

Most children can safely get all of these vaccines. But there are some exceptions:

- A child who has a mild cold or other illness on the day vaccinations are scheduled may be vaccinated. A child who is moderately or severely ill on the day of vaccinations might be asked to come back for them at a later date.
- Any child who had a life-threatening allergic reaction after getting a vaccine should not get another dose of that vaccine. **Tell the person giving the vaccines if your child has ever had a severe reaction after any vaccination.**
- A child who has a severe (life-threatening) allergy to a substance should not get a vaccine that contains that substance. **Tell the person giving your child the vaccines if your child has any severe allergies that you are aware of.**

Talk to your doctor before your child gets:

- **DTaP vaccine**, if your child ever had any of these reactions after a previous dose of DTaP:
 - A brain or nervous system disease within 7 days,
 - Non-stop crying for 3 hours or more,
 - A seizure or collapse,
 - A fever of over 105°F.
- **PCV13 vaccine**, if your child ever had a severe reaction after a dose of DTaP (or other vaccine containing diphtheria toxoid), or after a dose of PCV7, an earlier pneumococcal vaccine.

3 Risks of a Vaccine Reaction

With any medicine, including vaccines, there is a chance of side effects. These are usually mild and go away on their own. Most vaccine reactions are not serious: tenderness, redness, or swelling where the shot was given; or a mild fever. These happen soon after the shot is given and go away within a day or two. They happen with up to about half of vaccinations, depending on the vaccine.

Serious reactions are also possible but are rare.

Polio, Hepatitis B and Hib Vaccines have been associated only with mild reactions.

DTaP and Pneumococcal vaccines have also been associated with other problems:

DTaP Vaccine

- **Mild Problems:** Fussiness (up to 1 child in 3); tiredness or loss of appetite (up to 1 child in 10); vomiting (up to 1 child in 50); swelling of the entire arm or leg for 1-7 days (up to 1 child in 30)—usually after the 4th or 5th dose.
- **Moderate Problems:** Seizure (1 child in 14,000); non-stop crying for 3 hours or longer (up to 1 child in 1,000); fever over 105°F (1 child in 16,000).
- **Serious problems:** Long term seizures, coma, lowered consciousness, and permanent brain damage have been reported following DTaP vaccination. These reports are extremely rare.

Pneumococcal Vaccine

- **Mild Problems:** Drowsiness or temporary loss of appetite (about 1 child in 2 or 3); fussiness (about 8 children in 10).
- **Moderate Problems:** Fever over 102.2°F (about 1 child in 20).

After any vaccine:

Any medication can cause a severe allergic reaction. Such reactions from a vaccine are very rare, estimated at about 1 in a million doses, and would happen within a few minutes to a few hours after the vaccination.

As with any medicine, there is a very remote chance of a vaccine causing a serious injury or death.

The safety of vaccines is always being monitored. For more information, visit: www.cdc.gov/vaccinesafety/

4

What if there is a serious reaction?

What should I look for?

- Look for anything that concerns you, such as signs of a severe allergic reaction, very high fever, or unusual behavior.

Signs of a severe allergic reaction can include hives, swelling of the face and throat, and difficulty breathing. In infants, signs of an allergic reaction might also include fever, sleepiness, and disinterest in eating. In older children signs might include a fast heartbeat, dizziness, and weakness. These would usually start a few minutes to a few hours after the vaccination.

What should I do?

- If you think it is a severe allergic reaction or other emergency that can't wait, call 9-1-1 or get the person to the nearest hospital. Otherwise, call your doctor.

Afterward, the reaction should be reported to the Vaccine Adverse Event Reporting System (VAERS). Your doctor should file this report, or you can do it yourself through the VAERS web site at www.vaers.hhs.gov, or by calling **1-800-822-7967**.

VAERS does not give medical advice.

5

The National Vaccine Injury Compensation Program

The National Vaccine Injury Compensation Program (VICP) is a federal program that was created to compensate people who may have been injured by certain vaccines.

Persons who believe they may have been injured by a vaccine can learn about the program and about filing a claim by calling **1-800-338-2382** or visiting the VICP website at www.hrsa.gov/vaccinecompensation. There is a time limit to file a claim for compensation.

6

How can I learn more?

- Ask your healthcare provider. He or she can give you the vaccine package insert or suggest other sources of information.
- Call your local or state health department.
- Contact the Centers for Disease Control and Prevention (CDC):
 - Call **1-800-232-4636 (1-800-CDC-INFO)**
 - Visit CDC's website at www.cdc.gov/vaccines or www.cdc.gov/hepatitis

Vaccine Information Statement **Multi Pediatric Vaccines**

11/05/2015

42 U.S.C. § 300aa-26

Office Use Only



Pneumococcal Conjugate Vaccine (PCV13): *What You Need to Know*

Many Vaccine Information Statements are available in Spanish and other languages. See www.immunize.org/vis

Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite www.immunize.org/vis

1 Why get vaccinated?

Pneumococcal conjugate vaccine (PCV13) can prevent **pneumococcal disease**.

Pneumococcal disease refers to any illness caused by pneumococcal bacteria. These bacteria can cause many types of illnesses, including pneumonia, which is an infection of the lungs. Pneumococcal bacteria are one of the most common causes of pneumonia.

Besides pneumonia, pneumococcal bacteria can also cause:

- Ear infections
- Sinus infections
- Meningitis (infection of the tissue covering the brain and spinal cord)
- Bacteremia (bloodstream infection)

Anyone can get pneumococcal disease, but children under 2 years of age, people with certain medical conditions, adults 65 years or older, and cigarette smokers are at the highest risk.

Most pneumococcal infections are mild. However, some can result in long-term problems, such as brain damage or hearing loss. Meningitis, bacteremia, and pneumonia caused by pneumococcal disease can be fatal.

2 PCV13

PCV13 protects against 13 types of bacteria that cause pneumococcal disease.

Infants and young children usually need 4 doses of pneumococcal conjugate vaccine, at 2, 4, 6, and 12–15 months of age. In some cases, a child might need fewer than 4 doses to complete PCV13 vaccination.

A dose of PCV13 vaccine is also recommended for anyone **2 years or older** with certain medical conditions if they did not already receive PCV13.

This vaccine may be given to **adults 65 years or older** based on discussions between the patient and health care provider.

3 Talk with your health care provider

Tell your vaccine provider if the person getting the vaccine:

- Has had an **allergic reaction after a previous dose of PCV13, to an earlier pneumococcal conjugate vaccine known as PCV7, or to any vaccine containing diphtheria toxoid** (for example, DTaP), or has any **severe, life-threatening allergies**.
- In some cases, your health care provider may decide to postpone PCV13 vaccination to a future visit.

People with minor illnesses, such as a cold, may be vaccinated. People who are moderately or severely ill should usually wait until they recover before getting PCV13.

Your health care provider can give you more information.



4 Risks of a vaccine reaction

- Redness, swelling, pain, or tenderness where the shot is given, and fever, loss of appetite, fussiness (irritability), feeling tired, headache, and chills can happen after PCV13.

Young children may be at increased risk for seizures caused by fever after PCV13 if it is administered at the same time as inactivated influenza vaccine. Ask your health care provider for more information.

People sometimes faint after medical procedures, including vaccination. Tell your provider if you feel dizzy or have vision changes or ringing in the ears.

As with any medicine, there is a very remote chance of a vaccine causing a severe allergic reaction, other serious injury, or death.

5 What if there is a serious problem?

An allergic reaction could occur after the vaccinated person leaves the clinic. If you see signs of a severe allergic reaction (hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, or weakness), call **9-1-1** and get the person to the nearest hospital.

For other signs that concern you, call your health care provider.

Adverse reactions should be reported to the Vaccine Adverse Event Reporting System (VAERS). Your health care provider will usually file this report, or you can do it yourself. Visit the VAERS website at www.vaers.hhs.gov or call **1-800-822-7967**. *VAERS is only for reporting reactions, and VAERS staff do not give medical advice.*

6 The National Vaccine Injury Compensation Program

The National Vaccine Injury Compensation Program (VICP) is a federal program that was created to compensate people who may have been injured by certain vaccines. Visit the VICP website at www.hrsa.gov/vaccinecompensation or call **1-800-338-2382** to learn about the program and about filing a claim. There is a time limit to file a claim for compensation.

7 How can I learn more?

- Ask your health care provider.
- Call your local or state health department.
- Contact the Centers for Disease Control and Prevention (CDC):
 - Call **1-800-232-4636 (1-800-CDC-INFO)** or
 - Visit CDC's website at www.cdc.gov/vaccines



Rotavirus Vaccine: What You Need to Know

Many Vaccine Information Statements are available in Spanish and other languages. See www.immunize.org/vis

Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite www.immunize.org/vis

1 Why get vaccinated?

Rotavirus vaccine can prevent rotavirus disease.

Rotavirus causes diarrhea, mostly in babies and young children. The diarrhea can be severe, and lead to dehydration. Vomiting and fever are also common in babies with rotavirus.

2 Rotavirus vaccine

Rotavirus vaccine is administered by putting drops in the child's mouth. Babies should get 2 or 3 doses of rotavirus vaccine, depending on the brand of vaccine used.

- The first dose must be administered before 15 weeks of age.
- The last dose must be administered by 8 months of age.

Almost all babies who get rotavirus vaccine will be protected from severe rotavirus diarrhea.

Another virus called porcine circovirus (or parts of it) can be found in rotavirus vaccine. This virus does not infect people, and there is no known safety risk. For more information, see <http://wayback.archive-it.org/7993/20170406124518/https://www.fda.gov/BiologicsBloodVaccines/Vaccines/ApprovedProducts/ucm212140.htm>.

Rotavirus vaccine may be given at the same time as other vaccines.

3 Talk with your health care provider

Tell your vaccine provider if the person getting the vaccine:

- Has had an **allergic reaction after a previous dose of rotavirus vaccine**, or has any severe, life-threatening allergies.
- Has a **weakened immune system**.

- Has **severe combined immunodeficiency (SCID)**.
- Has had a type of bowel blockage called **intussusception**.

In some cases, your child's health care provider may decide to postpone rotavirus vaccination to a future visit.

Infants with minor illnesses, such as a cold, may be vaccinated. Infants who are moderately or severely ill should usually wait until they recover before getting rotavirus vaccine.

Your child's health care provider can give you more information.

4 Risks of a vaccine reaction

- Irritability or mild, temporary diarrhea or vomiting can happen after rotavirus vaccine.

Intussusception is a type of bowel blockage that is treated in a hospital and could require surgery. It happens naturally in some infants every year in the United States, and usually there is no known reason for it. There is also a small risk of intussusception from rotavirus vaccination, usually within a week after the first or second vaccine dose. This additional risk is estimated to range from about 1 in 20,000 US infants to 1 in 100,000 US infants who get rotavirus vaccine. Your health care provider can give you more information.

As with any medicine, there is a very remote chance of a vaccine causing a severe allergic reaction, other serious injury, or death.



5**What if there is a serious problem?**

For intussusception, look for signs of stomach pain along with severe crying. Early on, these episodes could last just a few minutes and come and go several times in an hour. Babies might pull their legs up to their chest. Your baby might also vomit several times or have blood in the stool, or could appear weak or very irritable. These signs would usually happen during the first week after the first or second dose of rotavirus vaccine, but look for them any time after vaccination. If you think your baby has intussusception, contact a health care provider right away. If you can't reach your health care provider, take your baby to a hospital. Tell them when your baby got rotavirus vaccine.

An allergic reaction could occur after the vaccinated person leaves the clinic. If you see signs of a severe allergic reaction (hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, or weakness), call **9-1-1** and get the person to the nearest hospital.

For other signs that concern you, call your health care provider.

Adverse reactions should be reported to the Vaccine Adverse Event Reporting System (VAERS). Your health care provider will usually file this report, or you can do it yourself. Visit the VAERS website at www.vaers.hhs.gov or call **1-800-822-7967**. *VAERS is only for reporting reactions, and VAERS staff do not give medical advice.*

6**The National Vaccine Injury Compensation Program**

The National Vaccine Injury Compensation Program (VICP) is a federal program that was created to compensate people who may have been injured by certain vaccines. Visit the VICP website at www.hrsa.gov/vaccinecompensation or call **1-800-338-2382** to learn about the program and about filing a claim. There is a time limit to file a claim for compensation.

7**How can I learn more?**

- Ask your health care provider.
- Call your local or state health department.
- Contact the Centers for Disease Control and Prevention (CDC):
 - Call **1-800-232-4636 (1-800-CDC-INFO)** or
 - Visit CDC's website at www.cdc.gov/vaccines

